



## Contains Non-Plant Food Ingredients

### Active Ingredients:

Blend of Nonionic Alkyl Glucamides and

Alkoxyated Polyols ..... **0.62%**

Inert Ingredients ..... **0.19%**

### Soil Amending Ingredients:

Bio-Mas Turf Enhancer ..... **0.03%**

Bio-Fusion Myco Rhizosphere Microbes ..... **0.05%**

Other Ingredients ..... **99.92%**

Rhizobial Microorganisms · *Bacillus subtilis* · *Bacillus/icheniformis* · *Bacillus megaterium* · *Bacillus pumilus* · *Bacillus azotoformans* · *Bacillus coagulans* · *Bacillus firmus* · *Bacillus amy/oliquefaciens* · *Paenibacillus polymyxa* · *Paenibacillus durum* · *Pseudomonas aureofaceans* · *Pseudomonas fluorescence* · *Streptomyces/ydicus* · *Streptomyces griseus* · *Trichoderma reesei* · *Trichoderma hamatum* · *Saccharomyces cerevisiae* · Endomycorrhizal Fungi: *G/omus aggregatum* · *G/omus mosseae* · *G/omus intraradices* Ectomycorrhizal Fungi: *Rhizopogon villosu/us* · *Rhizopogon luteo/us* · *Rhizopogon amy/opogon* · *Rhizopogon fulvigleba/solitus* *Ilncorius*

**Net Weight: 50 lbs. (18.1 kg)**



**ORGANIC  
PRECISE**<sup>TM</sup>

Powered by EBS<sup>o</sup>

### CAUTION: KEEP OUT OF REACH OF CHILDREN

WARNING: Causes mild skin irritation. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe dust. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves/protective clothing/eye protection/face protection. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. Rinse mouth. IF ON SKIN: Wash with plenty of water. Take off contaminated clothing and wash it before reuse. Get medical advice/attention if you feel unwell. Store locked up. Dispose of contents/container according to local regulations.

## Application Rates:

Turf Applications:

Application rate 4 lbs per 1,000 sq. ft. Apply at anytime during growing season.

Apply at 10,000 sq. ft. per bag (4 lbs per 1,000 sq. ft.

Ornamental Applications:

2-4 Tbsp/1 gal container, 2-4 Tbsp/1" caliber tree, 1-2 bags per 4 yards of media.

Annuals: 2-4 lbs per 1,000 sq. ft.

Directions for Use: Apply only as directed.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to, a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan. Some applications may require higher use rates and/or more frequent applications to produce significant benefits.

For Best Results water in with a minimum of 0.25 inches of water after application to incorporate into soil within 48 hours.

### PRECAUTIONARY STATEMENTS

Avoid prolonged or repeated contact with eyes, skin, and clothing. Chemical goggles or a full face shield should be worn. To protect skin wear appropriate protective equipment, such as rubber or plastic aprons, rubber gloves, and boots. Avoid breathing mist or vapor. Keep containers closed. Wash thoroughly after handling. May cause gastrointestinal distress if swallowed.

**Guaranteed by:**

**AUGI**

**517 Franklinville Rd.  
Mullica Hill, NJ 08062**



**Made In The U.S.A.**